

WARDOWSKI IMPLICIT CONTRACTIONS IN METRIC SPACES

MIHAI TURINICI

ABSTRACT. Most of the implicit contractions introduced by Wardowski [Fixed Point Th. Appl., 2012, 2012:94] are Matkowski type contractions.

1. INTRODUCTION

Let X be a nonempty set; and $d : X \times X \rightarrow R_+ := [0, \infty[$ be a *metric* over it; supposed to be *complete* [each d -Cauchy sequence is d -convergent]. Further, take some $T \in \mathcal{F}(X)$. [Here, given the nonempty sets A and B , $\mathcal{F}(A, B)$ stands for the class of all functions $f : A \rightarrow B$; when $A = B$, we write $\mathcal{F}(A, A)$ as $\mathcal{F}(A)$]. Denote $\text{Fix}(T) := \{z \in X; z = Tz\}$; any point of it will be called *fixed* under T . These points are to be determined in the context below (cf. Rus [8, Ch 2, Sect 2.2]):

1a) Call $x \in X$, a *semi-Picard point* (modulo (d, T)), provided $(T^n x; n \geq 0)$ is d -semi-Cauchy: $d(T^n x, T^{n+1} x) \rightarrow 0$. If this happens for each $x \in X$, we say that T is a *semi-Picard operator* (modulo d); and if (in addition), $\text{Fix}(T)$ is a *singleton* $[z, w \in \text{Fix}(T) \implies z = w]$, T is called a *global semi-Picard operator* (modulo d)

1b) Call $x \in X$, a *Picard point* (modulo (d, T)), provided $(T^n x; n \geq 0)$ is d -Cauchy; hence, d -convergent. If this happens for each $x \in X$, we say that T is a *Picard operator* (modulo d); and if (in addition), $\text{Fix}(T)$ is singleton, then T is referred to as a *global Picard operator* (modulo d)

1c) Call $x \in X$, a *super-Picard point* (modulo (d, T)), provided it is a Picard point (modulo (d, T)) and $z := \lim_n T^n x$ belongs to $\text{Fix}(T)$. If this happens for each $x \in X$, we say that T is a *super-Picard operator* (modulo d); and if, in addition, $\text{Fix}(T)$ is a singleton then T is called a *global super-Picard operator* (modulo d).

Sufficient conditions for such properties will be stated in the class of "functional" metric contractions. Call T , (d, φ) -*contractive* (for some $\varphi \in \mathcal{F}(R_+)$), when

$$(a01) \quad d(Tx, Ty) \leq \varphi(d(x, y)), \quad \forall x, y \in X, x \neq y.$$

The functions to be considered here are as follows. Let us say that $\varphi \in \mathcal{F}(R_+)$ is *increasing*, in case $[t_1 \leq t_2 \implies \varphi(t_1) \leq \varphi(t_2)]$; the class of all these will be denoted as $\mathcal{F}(i)(R_+)$. Further, let us call $\varphi \in \mathcal{F}(i)(R_+)$, *regressive* in case $\varphi(0) = 0$ and $[\varphi(t) < t, \forall t > 0]$. Finally, we shall say that $\varphi \in \mathcal{F}(i)(R_+)$ has the *Matkowski property* if $[\varphi^n(t) \rightarrow 0, \text{ for all } t > 0]$.

The following fixed point result in Matkowski [5] is our starting point.

Theorem 1. *Supposed that T is (d, φ) -contractive, for some regressive Matkowski function $\varphi \in \mathcal{F}(i)(R_+)$. Then, T is a global super-Picard operator.*

2010 *Mathematics Subject Classification.* 47H10 (Primary), 54H25 (Secondary).

Key words and phrases. Metric space, Picard operator, fixed point, regressive Matkowski function, Wardowski contraction.

In particular, when φ is *linear* (i.e.: $\varphi(t) = \alpha t$, $t \in R_+$, for some $\alpha \in]0, 1[$) Theorem 1 is just the 1922 Banach's contraction mapping principle [3]. This result found some interesting applications in operator equations theory; so, it was the subject of various extensions. For example, a natural way of doing this consists in taking implicit "functional" contractive conditions like

$$(a02) \quad F(d(Tx, Ty), d(x, y)) \leq 0, \quad \forall x, y \in X, x \neq y;$$

where $F : R_+^2 \rightarrow R \cup \{-\infty, \infty\}$ is an appropriate function. For more details about the possible choices of F we refer to the 1976 paper by Turinici [9]. A further extension of this is by considering implicit functional contractive conditions like

$$(a03) \quad F(d(Tx, Ty), d(x, y), d(x, Tx), d(y, Ty), d(x, Ty), d(y, Tx)) \leq 0, \\ \forall x, y \in X, x \neq y;$$

where $F : R_+^6 \rightarrow R \cup \{-\infty, \infty\}$ is an appropriate function. Some concrete examples may be found in Berinde and Vetro [4]; see also Rhoades [7] and Akkouchi [2].

Recently, an interesting contractive condition of the type (a02) was introduced in Wardowski [11]. It is our aim in the following to show that, concerning the fixed point question, a reduction to Theorem 1 is possible for most of these contractions; cf. Section 3. For the remaining ones, we provide in Section 4 a result where some specific requirements posed by Wardowski (cf. Section 5) are shown to be superfluous. Finally, Section 2 has an introductory character. Further aspects will be delineated in a future paper.

2. PRELIMINARIES

(A) Let in the following $R_+^0 :=]0, \infty[$ stand for the subset of all strict positive (real) numbers. Call $F : R_+ \rightarrow R \cup \{-\infty\}$, a *semi-Wardowski* function, provided

$$(b01) \quad F(t) = -\infty \text{ if and only if } t = 0$$

$$(b02) \quad F \text{ is strictly increasing: } t < s \implies F(t) < F(s).$$

As a consequence of these facts, the lateral limits

$$F(t-0) := \lim_{s \rightarrow t-} F(s), \quad F(t+0) := \lim_{s \rightarrow t+} F(s) \quad (2.1)$$

exist, for each $t > 0$; in addition,

$$-\infty < F(t-0) \leq F(t) \leq F(t+0) < \infty, \quad \forall t > 0. \quad (2.2)$$

Note that, in general, F is not continuous. However, by the general properties of the monotone functions, we have (cf. Natanson, [6, Ch 8, Sect 1]):

Proposition 1. *Suppose that $F : R_+ \rightarrow R \cup \{-\infty\}$ is a semi-Wardowski function. Then, there exists a countable subset $\Delta(F) \subseteq R_+^0$ such that*

$$F(t-0) = F(t) = F(t+0), \text{ for each } t \in R_+^0 \setminus \Delta(F). \quad (2.3)$$

In addition, the following property of such objects is useful for us.

Lemma 1. *Let $F : R_+ \rightarrow R \cup \{-\infty\}$ be a semi-Wardowski function. Then,*

$$\forall t, s \in R_+ : F(t) < F(s) \implies t < s. \quad (2.4)$$

Proof. Take the numbers $t, s \in R_+$ according to the premise of this relation. Clearly, $s > 0$; and, from this, the case $t = 0$ is proved; hence, we may assume that $t > 0$. The alternative $t = s$ gives $F(t) = F(s)$; in contradiction with $F(t) < F(s)$. Likewise, the alternative $t > s$ yields (via (b02)) $F(t) > F(s)$; again in contradiction with $F(t) < F(s)$. Hence, $t < s$; and the conclusion follows. $\square$

(B) Now, let us add one more condition (upon such functions)

(b03) $F(t) \rightarrow -\infty$, when $t \rightarrow 0+$; i.e.: $F(0+) = -\infty$.

When F satisfies (b01)-(b03), it will be referred to as a *Wardowski* function.

Lemma 2. *Suppose that $F : R_+ \rightarrow R \cup \{-\infty\}$ is a Wardowski function. Then, for each sequence (t_n) in R_+^0 ,*

$$F(t_n) \rightarrow -\infty \text{ implies } t_n \rightarrow 0. \quad (2.5)$$

Proof. Suppose that this is not true: there must be some $\varepsilon > 0$ such that

for each n , there exists $m > n$, such that $t_m \geq \varepsilon$.

We get therefore a subsequence $(s_n := t_{i(n)})$ of (t_n) such that

$$s_n \geq \varepsilon \text{ (hence, } F(s_n) \geq F(\varepsilon)), \forall n.$$

This, however, contradicts the property $F(s_n) \rightarrow -\infty$; hence the conclusion. $\square$

3. LEFT-CONTINUOUS WARDOWSKI FUNCTIONS

(A) Suppose that $F : R_+ \rightarrow R \cup \{-\infty\}$ is a Wardowski function; and let $a > 0$ be fixed in the sequel. Denote, for each $t \geq 0$,

$$(c01) \quad M(a, F)(t) = \{s \geq 0; a + F(s) \leq F(t)\}, \quad \varphi(t) = \sup M(a, F)(t).$$

Note that, from (b03), $M(a, F)(t)$ is nonempty, for each $t \geq 0$; and $M(a, F)(0) = \{0\}$ [hence $\varphi(0) = 0$]. In addition, for each $t > 0$ one has, via Lemma 1,

$$s \in M(a, F)(t) \implies F(s) < F(t) \implies s < t;$$

wherefrom, again for all $t > 0$,

$$M(a, F)(t) \subseteq [0, t]; \text{ whence, } \varphi(t) \leq t. \quad (3.1)$$

Moreover, as F is (strictly) increasing,

$$0 \leq t_1 \leq t_2 \implies M(a, F)(t_1) \subseteq M(a, F)(t_2) \implies \varphi(t_1) \leq \varphi(t_2); \quad (3.2)$$

so that, φ is increasing on R_+ .

A basic problem to be posed is that of determining sufficient conditions under F such that the (increasing) function φ be regressive Matkowski (cf. Section 1). The following answer is available.

Proposition 2. *Suppose that the Wardowski function $F : R_+ \rightarrow R \cup \{-\infty\}$ fulfills*

$$(c02) \quad F \text{ is left-continuous: } F(t-0) = F(t), \forall t > 0.$$

Then, the associated (increasing) function φ is regressive and Matkowski.

Proof. There are two steps to be passed.

Step 1. Let $t > 0$ be arbitrary fixed; and put $u := \varphi(t)$. If $u = 0$, we are done; so, without loss, we may assume that $u > 0$. By the very definition of this number,

$$a + F(s) \leq F(t), \text{ for all } s \in [0, u].$$

So, passing to limit as $(s \rightarrow u-)$, one gets, via (c02),

$$a + F(u) \leq F(t); \text{ i.e.: } a + F(\varphi(t)) \leq F(t). \quad (3.3)$$

This in turn yields $F(\varphi(t)) < F(t)$; whence $\varphi(t) < t$. As $t > 0$ was arbitrarily chosen, it follows that φ is regressive.

Step 2. Fix some $t > 0$; and let the iterative sequence (t_n) be given as $[t_0 = t, t_{n+1} = \varphi(t_n); n \geq 0]$. If $t_h = 0$, for some index $h > 0$, we are done; so, without loss, one may assume that $[t_n > 0, \forall n]$. By (3.3), we have

$$a + F(t_{n+1}) \leq F(t_n), [\text{hence, } a \leq F(t_n) - F(t_{n+1})], \forall n.$$

Adding the first n inequalities, one gets

$$na \leq F(t_0) - F(t_n) [\text{hence: } F(t_n) \leq F(t_0) - na], \forall n;$$

so that, passing to limit as $n \rightarrow \infty$, one derives $F(t_n) \rightarrow -\infty$. This, along with Lemma 2, gives $t_n \rightarrow 0$. As $t > 0$ was arbitrarily chosen, it results that φ has the Matkowski property. The proof is finished. $\square$

(B) Let (X, d) be a complete metric space; and $T \in \mathcal{F}(X)$, a selfmap of X . Given the real number $a > 0$ and the Wardowski function $F : R_+ \rightarrow R \cup \{-\infty\}$, let us say that T is (a, F) -contractive, if

$$(c03) \ a + F(d(Tx, Ty)) \leq F(d(x, y)), \forall x, y \in X, x \neq y.$$

The first main result of this exposition is

Theorem 2. *Suppose that T is (a, F) -contractive, for some $a > 0$ and some left-continuous Wardowski function $F(\cdot)$. Then, T is a global super-Picard map.*

Proof. By Proposition 2, the associated by (c01) increasing function $\varphi \in \mathcal{F}(R_+)$ is regressive Matkowski. On the other hand, (c03) tells us that T is (d, φ) -contractive. This, along with Theorem 1, gives us all conclusions we need. $\square$

(C) Given $k > 0$, let us say that the Wardowski function $F : R_+ \rightarrow R \cup \{-\infty\}$ is k -regular, provided $[\lim_{t \rightarrow 0+} t^k F(t) = 0]$. As a direct consequence of Theorem 2 we have, formally,

Theorem 3. *Suppose that T is (a, F) -contractive, for some $a > 0$ and some left-continuous Wardowski function $F : R_+ \rightarrow R \cup \{-\infty\}$, with*

$$(c04) \ F \text{ is } k\text{-regular, for some } k \in]0, 1[.$$

Then, T is a global super-Picard map.

For a different way of proving this result we refer to Section 5 below. Note that all examples in Wardowski [11] are illustrations of Theorem 3; such as

$$(c05) \ F(0) = -\infty; F(t) = \log(\alpha t^2 + \beta t) + \gamma t, t > 0,$$

$$(c06) \ F(0) = -\infty; F(t) = (-1)(t^{-\delta}), t > 0;$$

here, $\log$ is the natural logarithm and $\alpha, \beta, \gamma > 0, \delta \in]0, 1[$ are constants. The case $\delta = 1$ of the last example is not reducible to Theorem 3. However, it is possible to handle it with the aid of Theorem 2; we do not give details.

4. DISCONTINUOUS WARDOWSKI FUNCTIONS

Let us now return to the general setting above. As results from Proposition 2, the left continuity requirement (c02) is essential in Theorem 2 so as to deduce it from Theorem 1. In the absence of this, the reduction argument above does not work. And then, the question arises of to what extent is Theorem 2 retainable in such an extended setting. Strange enough, a positive answer to this is still available. But, before stating it, some preliminaries are needed.

(A) Let (X, d) be a complete metric space. Remember (cf. Section 1) that a sequence $(x_n; n \geq 0)$ in X is called d -semi-Cauchy, provided $d(x_n, x_{n+1}) \rightarrow 0$.

In the following, a useful property is described for such sequences which are not d -Cauchy. For the sequence $(r_n; n \geq 0)$ in R_+^0 and the point $r \in R_+^0$, let us write $r_n \downarrow r$ when $[r_n > r, \forall n]$ and $r_n \rightarrow r$.

Proposition 3. *Suppose that $(x_n; n \geq 0)$ is d -semi-Cauchy but not d -Cauchy. Further, let Δ be a countable part of R_+^0 . There exist then a number $\eta \in R_+^0 \setminus \Delta$, a rank $j(\eta) \geq 0$, and a couple of rank-sequences $(m(j); j \geq 0)$, $(n(j); j \geq 0)$ with*

$$j \leq m(j) < n(j), \quad d(x_{m(j)}, x_{n(j)}) > \eta, \quad \forall j \geq 0 \quad (4.1)$$

$$n(j) - m(j) \geq 2, \quad d(x_{m(j)}, x_{n(j)-1}) \leq \eta, \quad \forall j \geq j(\eta) \quad (4.2)$$

$$(d(x_{m(j)}, x_{n(j)}); j \geq 0) \downarrow \eta \quad (4.3)$$

$$(d(x_{m(j)+p}, x_{n(j)+q}); j \geq 0) \rightarrow \eta, \quad \forall p, q \in \{0, 1\}. \quad (4.4)$$

Proof. By definition, the d -Cauchy property of our sequence means:

$$\forall \varepsilon \in R_+^0, \exists k = k(\varepsilon): k \leq m \leq n \implies d(x_m, x_n) \leq \varepsilon.$$

As $R_+^0 \setminus \Delta$ is dense in R_+^0 , this property may be also written as

$$\forall \eta \in R_+^0 \setminus \Delta, \exists k = k(\eta): k \leq m \leq n \implies d(x_m, x_n) \leq \eta.$$

The negation of this property means that, there must be some $\eta \in R_+^0 \setminus \Delta$ in such a way that, $\forall j \geq 0$:

$$A(j) := \{(m, n) \in N \times N; j \leq m < n, d(x_m, x_n) > \eta\} \neq \emptyset. \quad (4.5)$$

Having this precise, denote, for each $j \geq 0$,

$$(d01) \quad m(j) = \min \text{Dom}(A(j)), \quad n(j) = \min A(m(j)).$$

As a consequence, the couple of rank-sequences $(m(j); j \geq 0)$, $(n(j); j \geq 0)$ fulfills (4.1). On the other hand, letting the index $j(\eta)$ be such that

$$\rho_i := d(x_i, x_{i+1}) < \eta, \quad \text{for all } i \geq j(\eta), \quad (4.6)$$

it is clear that (4.2) holds too. This in turn yields (by the triangular inequality)

$$\eta < d(x_{m(j)}, x_{n(j)}) \leq d(x_{m(j)}, x_{n(j)-1}) + \rho_{n(j)-1} \leq \eta + \rho_{n(j)-1}, \quad \forall j \geq j(\eta);$$

so, passing to limit as $j \rightarrow \infty$ gives (4.3). Finally, again by the triangular inequality, one has, for all $j \geq 0$,

$$d(x_{m(j)}, x_{n(j)+1}) \leq d(x_{m(j)}, x_{n(j)}) + \rho_{n(j)},$$

$$d(x_{m(j)}, x_{n(j)+1}) \geq d(x_{m(j)}, x_{n(j)}) - \rho_{n(j)}.$$

By a limit process upon j one gets the case $(p = 0, q = 1)$ of (4.4). The remaining ones are obtained in a similar way. $\square$

(B) We are now in position to state the second main result of this exposition. Let $T \in \mathcal{F}(X)$ be a selfmap of X .

Theorem 4. *Suppose that T is (a, F) -contractive, for some $a > 0$ and some Wardowski function $F : R_+ \rightarrow R \cup \{-\infty\}$. Then, T is a global super-Picard map.*

Proof. From the (a, F) -contractive condition, it results that

$$T \text{ is a strict contraction: } d(Tx, Ty) < d(x, y), \quad \forall x, y \in X, x \neq y. \quad (4.7)$$

This firstly gives us that $\text{Fix}(T)$ is a singleton. As a second consequence,

$$T \text{ is nonexpansive: } d(Tx, Ty) \leq d(x, y), \quad \forall x, y \in X; \quad (4.8)$$

hence, in particular, T is d -continuous on X . So, it remains only to prove that T is a Picard operator. Let $x = x_0$ be arbitrary fixed in X ; and put $(x_n := T^n x; n \geq 0)$. If $x_k = x_{k+1}$ for some $k \geq 0$, we are done; so, without loss, one may assume that

$$(d02) \quad \rho_n := d(x_n, x_{n+1}) > 0, \forall n.$$

Part 1. By the contractive condition, we have

$$a + F(\rho_{n+1}) \leq F(\rho_n), [\text{hence, } a \leq F(\rho_n) - F(\rho_{n+1})], \forall n.$$

Adding the first n inequalities, one gets

$$na \leq F(\rho_0) - F(\rho_n) [\text{hence: } F(\rho_n) \leq F(\rho_0) - na], \forall n;$$

so that, passing to limit as $n \rightarrow \infty$, one derives $F(\rho_n) \rightarrow -\infty$. This, along with Lemma 2, gives $\rho_n \rightarrow 0$; hence, $(x_n; n \geq 0)$ is d -semi-Cauchy.

Part 2. Let $\Delta(F)$ stand for the subset of R_+^0 where F is discontinuous; note that, by Proposition 1, it is (at most) countable. Assume by contradiction that (x_n) is not d -Cauchy. By Proposition 3, there exists a number $\eta \in R_+^0 \setminus \Delta(F)$, a rank $j(\eta) \geq 0$, and a couple of rank-sequences $(m(j); j \geq 0)$, $(n(j); j \geq 0)$ with the properties (4.1)-(4.4). For simplicity, we shall write (m, n) in place of $(m(j), n(j))$, respectively. By the contractive condition, we have

$$a + F(d(x_{m+1}, x_{n+1})) \leq F(d(x_m, x_n)), \forall j.$$

Passing to limit as $j \rightarrow \infty$ one gets, from the choice of η and (4.3)-(4.4),

$$a + F(\eta) \leq F(\eta); \text{ hence, } a \leq 0.$$

The obtained contradiction tells us that (x_n) is d -Cauchy; and, from this, all is clear via (4.8). The proof is complete. $\square$

5. FURTHER ASPECTS

Let again (X, d) be a complete metric space; and $T \in \mathcal{F}(X)$ be a selfmap of X . From Theorem 4, we have formally

Theorem 5. *Suppose that T is (a, F) -contractive, for some $a > 0$ and some Wardowski function $F(\cdot)$, taken as in (c04). Then, T is a global super-Picard map.*

This result is, essentially, the one in Wardowski [11]. For completeness reasons, we shall provide its proof, with some modifications.

Proof. (Theorem 5) As in Theorem 4, we only have to establish that T is a Picard operator. Let $x = x_0$ be arbitrary fixed in X ; and put $[x_n := T^n x, n \geq 0]$. If $x_k = x_{k+1}$ for some $k \geq 0$, we are done; so, without loss, one may assume that (d02) holds. By the contractive condition, we have

$$a + F(\rho_{n+1}) \leq F(\rho_n), [\text{hence, } a \leq F(\rho_n) - F(\rho_{n+1})], \forall n.$$

Adding the first n inequalities, one gets

$$na \leq F(\rho_0) - F(\rho_n) [\text{hence: } F(\rho_n) \leq F(\rho_0) - na], \forall n; \quad (5.1)$$

This firstly gives (passing to limit as $n \rightarrow \infty$), $F(\rho_n) \rightarrow -\infty$; wherefrom, by Lemma 2, $\rho_n \rightarrow 0$; hence, $(x_n; n \geq 0)$ is d -semi-Cauchy. Secondly, (5.1) also gives

$$na\rho_n^k \leq [F(\rho_0) - F(\rho_n)]\rho_n^k, \forall n. \quad (5.2)$$

By the deduced property of $(\rho_n; n \geq 0)$ and (c04), the limit of the right hand side is zero; so that, given $\beta > 0$, there must be some rank $i = i(\beta)$ such that

$$[F(\rho_0) - F(\rho_n)]\rho_n^k \leq \beta, \quad \forall n \geq i.$$

Combining with (5.2) yields (after transformations)

$$\rho_n \leq (\beta/an)^{1/k}, \text{ for all } n \geq i.$$

This, along with the convergence of the series $\sum_{n \geq 1} n^{-1/k}$ tells us that so is the series $\sum_n \rho_n$; wherefrom, $(x_n; n \geq 0)$ is d -Cauchy. The last part is identical with the one of Theorem 4; and conclusion follows. $\square$

Note, finally, that all these results are extendable to the framework of quasi-ordered metric spaces under the lines in Agarwal et al [1]; see also Turinici [10]. A development of these facts will be given in a future paper.

REFERENCES

- [1] R. P. Agarwal, M. A. El-Gebeily and D. O'Regan, *Generalized contractions in partially ordered metric spaces*, Appl. Anal., 87 (2008), 109-116.
- [2] M. Akkouchi, *Coincidence points for multivalued mappings*, An. Șt. Univ. Ovidius Constanța, 18 (2010), 13-22.
- [3] S. Banach, *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fund. Math., 3 (1922), 133-181.
- [4] V. Berinde and F. Vetro, *Common fixed points of mappings satisfying implicit contractive conditions*, Fixed Point Th. Appl., 2012, 2012:105.
- [5] J. Matkowski, *Integrable solutions of functional equations*, Dissertationes Math., 127 (1975), 1-68.
- [6] I. P. Natanson, *Theory of Real Variable Functions* (Russian), Gos. Izd. Teh.-Teor. Lit., Moskva, 1950.
- [7] B. E. Rhoades, *A comparison of various definitions of contractive mappings*, Trans. Amer. Math. Soc., 226 (1977), 257-290.
- [8] I. A. Rus, *Generalized Contractions and Applications*, Cluj University Press, Cluj-Napoca, 2001.
- [9] M. Turinici, *Fixed points of implicit contraction mappings*, An. Șt. Univ. "Al. I. Cuza" Iași (S I-a, Mat), 22 (1976), 177-180.
- [10] M. Turinici, *Fixed points for monotone iteratively local contractions*, Dem. Math., 19 (1986), 171-180.
- [11] D. Wardowski, *Fixed points of a new type of contractive mappings in complete metric spaces*, Fixed Point Th. Appl., 2012, 2012:94.

"A. MYLLER" MATHEMATICAL SEMINAR; "A. I. CUZA" UNIVERSITY; 700506 IAȘI, ROMANIA
E-mail address: `mturi@uaic.ro`